

LA-UR-11-10660

Approved for public release; distribution is unlimited.

Title: MiniBooNE Neutrino Oscillation Results

Author(s): Louis, William C. III

Intended for: Poster for general viewing



Disclaimer:



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By acceptance of this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.




# Neutrino Oscillation Results from MiniBooNE

The MiniBooNE experiment has searched for neutrino oscillations with both a neutrino beam and an antineutrino beam. In neutrino mode, MiniBooNE observes no oscillations consistent with the signal observed by the LSND experiment (A. A. Aguilar-Arevalo et al., Phys. Rev. Lett. 102, 101802 (2009)). However, the antineutrino data are consistent with oscillations at  $\sim 1 \text{ eV}^2$  and consistent with the LSND signal (A. A. Aguilar-Arevalo et al., Phys. Rev. Lett. 105, 181801 (2010)). These data suggest the possibility of CP violation in the lepton sector.


A schematic drawing of the MiniBooNE experiment. Protons from the 8-GeV Booster accelerator at Fermilab interact in a beryllium target, producing pions that are focused by a magnetic focusing horn. The pions then decay in a 50 m decay pipe into muons and neutrinos. The MiniBooNE detector, located 541 m downstream of the beryllium target, consists of a 40-foot spherical tank filled with  $\sim 800$  tons of mineral oil and covered on the inside by 1520 8-inch photobes. The detector reconstructs neutrinos that interact in the mineral oil.




The excess number of events as a function of  $L/E$  for LSND antineutrino data (black points) and MiniBooNE antineutrino data (red points).  $L$  is the distance traveled by the antineutrinos, while  $E$  is the energy of the antineutrinos. The two distributions are consistent, as expected from oscillations.




The excess events observed in neutrino mode as a function of reconstructed neutrino energy. Although an excess of events is observed at low energies, no appreciable event excess is observed at higher energy.



The limit on oscillations observed in neutrino mode (black curve) compared with the oscillation region from the LSND experiment (blue region). The MiniBooNE neutrino data are not consistent with the LSND signal.



The excess events observed in antineutrino mode as a function of reconstructed antineutrino energy. Excess events are observed at both low and high energies.



The oscillation region in antineutrino mode (closed curves) compared with the oscillation region from the LSND experiment (blue region). The MiniBooNE data are consistent with the LSND signal in antineutrino mode.

LA-UR-11-XXXX  
Contact info or website?